Bermuda & Zoysia Lawn Care Made East

Turf & Step HiYield® Weed **Ornamental** & Grass Stopper Con-

Dimension may taining be applied in winter & again in spring for the prevention of many common lawn weeds. It will also stop crabgrass from germinating & destroys the weed during the first five weeks of its growth.



WEED FREE ZONE Containing Lawn

Fertilizer Should be applied early spring through May. Wet the lawn be-

fore applying, wait 24 hours then water thoroughly. This product will control most common broadleaf weeds while providing a balanced lawn food. For more thor-



ough weed control spray ferti-lome® Weed Free Zone 10-14 days later.

> Winterizer for Es-

Lawn Food Plus

Iron. Apply in June. This product will noticeably green-up a lawn almost "overnight" & keep it "green" long term. This lawn food contains Chelated

iron which is rapidly available to the turf. For best results apply monthly in the summer, weather permitting.

tablished Lawns Apply in the fall to both lawns & shrubs. It better enables turf, trees & shrubs to withstand the rigors of winter. Its effects are similar to putting on an overcoat in the winter.



Extra potash enables Warm Season

grasses over winter better and be healthier in the spring. For additional weed control apply ferti-lome® Winterizer & Weed Preventer II For Southern Grasses...





Winterizer

Apply ferti-lome® F-Stop For Picture Perfect Disease Free Turfgrass. In severe cases, repeated applications may be necessary. It is advisable to apply a preventive application of ferti-lome® F-Stop in the spring & fall. Specific types of crawling insects (such as ants, crickets, grasshoppers etc.) may be controlled by monthly applications of Hi-Yield® Bug Blaster II Turf Insect Control Granules.

awn Food

Plus Iron

GREENER!



All Seasons Mulch Market & Landscape Supply

1820 River Road | Johns Island, SC 29455 Phone: (843) 559-5829

Always read & follow label directions

